

Make Every Minute Count

AFINION™ 2 ANALYZER

The Afinion™ 2 Analyzer is the result of years of continuous product innovation and refinement and delivers actionable point-of-care test results in just minutes.



AFINION™ 2 ANALYZER

With its compact size and panel of tests, the AfinionTM 2 System is ideal for point-of-care testing in general practice, community pharmacy, diabetes clinics, community health centers, hospital emergency departments and pathology laboratories. From just a small urine or fingerstick whole blood sample, highly accurate results for CRP, HbA1c, ACR and Lipid Panel are made available during the consultation.

BENEFITS TO POINT-OF-CARE TESTING

Reduced workload in the practice1:

- · Fewer follow-up calls
- Fewer follow-up letters
- Fewer patient revisits

More informed decision-making improves patient-clinician relationships and patient motivation and attitude.²

TESTS

Afinion™ tests can be used with both the Afinion™ 2 Analyzer and the Afinion™ AS100 Analyzer.

PRODUCT NAME	PRODUCT CODE	TESTS/BOX
AFINION™ 2 ANALYZER	1116770	N/A
AFINION™ HbA1c	1116795	15
AFINION™ ACR	1116781	15
AFINION™ CRP	1116787	15
AFINION™ LIPID PANEL	1116801	15

TECHNICAL INFORMATION

- Size: 200 mm W × 186 mm H × 328 mm D
- Weight: 3.4 kg
- LCD color touch display
- Patient and QC ID and results stored in memory
- No maintenance required
- No operator calibration
- Extensive and comprehensive fail-safe systems prevent the possibility of erroneous results
- · QC and operator lockout

CONNECTIVITY

The Afinion 2 Analyzer can reliably transfer test results to an information system. Use the Ethernet® cable to interface the analyzer to an information system. The Afinion 2 Analyzer connectivity automatically transfers patient and control results to a connected LIS/HIS/EMR system via TCP/IP networking using the protocols POCT1-A, HL7®, ASTM

1381-95 (low level) or ASTM 1394-97 (high level), selectable by configuration. With built-in connectivity, the Afinion 2 Analyzer can communicate with LIS and HIS systems to share and manage data more effectively, contributing to better patient management everywhere.

3-STEP PROCEDURE

Collect the sample with the integrated sampling device.



Place the sampling device back in the test cartridge.



Place the test cartridge in the analyzer and close the lid. The processing starts automatically.



CONTACT US TODAY FOR MORE INFORMATION. TO ARRANGE A DEMONSTRATION, VISIT GLOBALPOINTOFCARE.ABBOTT

 $T: 1800\ 622\ 642\ (free\ call\ AU)\ |\ T: 0800\ 466\ 951\ (free\ call\ NZ)\ |\ T: +61\ 7\ 3363\ 7100\ |\ rapiddx. ANZ. enquiries@abbott.com$

- 1. Crocker JB, Lee-Lewandrowski E, Lewandroski N, et al. Implementation of point-of-care testing in an ambulatory practice of an academic medical center. Am J Clin Pathol. 2014;142:640-6.
- 2. Laurence CO, Gialamas A, Bubner T. Patient satisfaction with point-of-care testing in general practice. Br J Gen Pract. 2010;60(572):e98-e104.

© 2020 Abbott. All rights reserved. All trademarks referenced are trademarks of either the Abbott group of companies or their respective owners. Any photos displayed are for illustrative purposes only. Any person depicted in such photos is a model. A2909.5 12/20

